

Supporting Information

Polystyrene microparticles size-controlled by thermo-dynamic emulsion polymerization

Sudip Kumar Pal^{1#}, Tohid Otoufat^{2#}, Gunwoo Kim^{1*}

1. Department of Organic Materials and Textile Engineering, Jeonbuk National University, 567 Baekje-daero, Deokjin-gu, Jeonju-si, Jeonbuk 54896, Republic of Korea.

2. Biomedical Manufacturing Technology Center, Korea Institute of Industrial Technology, Yeongcheon 38822, Republic of Korea.

*Corresponding author. Email: creaventure@jbnu.ac.kr (G.K.)

Table of Content

Figure S1. IR spectrums of all synthesized polystyrene particles.

Figure S2. Topographic chart of size distribution with all the conditions of temperature and initiator concentration. (A) Size of the 1st decimal particle (B) Size of the 9th decimal particle.

Figure S3. Graph of statistical data of the size of synthesized polystyrene particles. (A) Solid concentration compares to Sphere particles yield (B) all conditions size distribution graph.

Figure S4. Graph of synthesized polystyrene particles solid concentration per time.

Figure S5. Size distribution diagrams of all centration conditions (A) [KPS] = 0.5% (B) [KPS] = 0.1% (C) [KPS] = 0.05%.




Figure 1. IR spectra of all synthesized polystyrene particles.




Figure S2. Topographic chart of size distribution with all the conditions of temperature and initiator concentration. (A) Size of the 1st decimal particle (B) Size of the 9th decimal particle.




Figure S3. Graph of statistical data of the size of synthesized polystyrene particles. (A) Solid concentration compares to Sphere particles yield (B) all conditions size distribution graph




Figure S4. Graph of synthesized polystyrene particles solid concentration per time.




Figure S5. Size distribution diagrams of all centration conditions (A) [KPS] = 0.5% (B) [KPS] = 0.1% (C) [KPS] = 0.05%